### Last time...

- What are monkey grade primates?
- When and where do they first appear in the fossil record?
- What features are found in the earliest representatives that identify them as monkeys?

### Questions from last class?

- Where do we find primates in the Oligocene?
- Of the Oligocene primates...
  - Who is the potential ancestor of the Catarrhines?
  - Who is the potential ancestor to the NWM?
    - How did the NWM get to South America?
  - Who is the potential ancestor to the OWM?

## Taxonomy


- How are monkeys classified?
  - Order? Suborder? Infraorder?
- What are the two groups of monkeys?
  - How do their classifications differ?
  - How do their characteristics differ?
- Of the Simiiformes, which two groups are most closely related?

# New World Monkeys

- What are the potential evolutionary sources of the New World Monkeys
- What is their classification?
- What features distinguish them from other primates and in particular from the Old World monkeys?
- What different kinds of monkeys are New World Monkeys?
- How do Callitrichidae differ from the rest of the Ceboidea?

# Old World Monkeys

- What are the potential evolutionary sources of the Old World Monkeys
- What is their classification?
- What features distinguish them from other primates and in particular from the New World monkeys?
- What different kinds of monkeys are Old World Monkeys?
- How do Cercopithecinae and Colobinae, the two divisions, differ from one another?



# Modern Apes















### **Living Primates**





Figure 5.5 Taxonomy of the Haplorrhines to the Family Level

### Ape Grade Characteristics


- No tail
- Y-5 Molars
- larger body
- larger brain
- greater social complexity
- shoulder joint adaptations




# Miocene 3.5-5 mya

- Ape grade
- Over 30 genera and 100 species of ape
  - compared with 6 today
- Africa and Eurasia


### Miocene continents






Note the prominent canine jugum (facial pillar) on the *Proconsul* skull, which is a trait *Proconsul* shares with other early hominoids, which holds the root of the upper canine. *Proconsul* is also characterized by a broad interorbital region and broad orbits.

Also notice how the upper central incisors are larger than the lateral ones, which is similar to later apes, and how narrow the lower incisors are.




### Juvenile Proconsul heseloni



Like quadrupedal primates, *Proconsul* has a narrow and deep thorax and a long flexible lumbar region. It likely had six lumbar vertebrae, similar to cercopithecoids and some gibbons, and lacked the specialized lumbar reduction and stiffening of the great apes.


While resembling quadrupedal monkeys in many postcranial features, *Proconsul* shares the derived loss of a tail with the extant apes, among other features.



Unlike later suspensory apes, *Proconsul* was an arboreal quadruped.


http://www.mc.maricopa.edu/~reffland/anthropology/anthro2003/origins/ primates/evolution.html

# Morotopithecus





### Dryopithecus




Dryopithecus shares nasal aperture and periorbital morphology with the African great apes. Also notice the narrow upper incisors, which are a derived trait unique to this genus.

Gorilla image source: http://www.amonline.net.au/human\_evolution/skulls/


# Oreopithecus






The cranium, while damaged and distorted in this image, has strong nuchal and sagittal crests which meet, smooth gracile orbital rims, a short snout, and a tubular ectotympanic.


The brain size is small compared to the other great apes, and is suggested to be a result of secondary reduction that is often associated with specialized forms of folivory.




Like the extant hominoids, *Oreopithecus* possesses a broad thorax, a short trunk, and evidence of extensive mobility in almost all joints. It is likely that *Oreopithecus* was highly suspensory.

### Pierolapithecus







### Reconstruction of Pierolapithecus









Notice how high the lower margin of the orbits are from top of the nasal aperture, which is a trait Sivapithecus shares with Pongo.





Whitehead et al. 2005

Sivapithecus and Pongo are similar in their possession of narrow interorbital regions, broad zygomatic arches, tall narrow nasal apertures, and high orbits, compared to Pan.




Whitehead et al. 2005

Notice how large and procumbent the premaxilla of Sivapithecus is compared to Pan, causing it to have a "dished" face. This is another derived trait Sivapithecus shares with Pongo.


# Gigantopithecus














# Superfamily: Hominoidea

- Gibbons, Gorillas, Orangutan, Chimpanzee, Human
- Greater encephalization, larger body, brachiation, social complexity, lack of tail

### Family: Hylobatidae - Gibbons



# Hylobatidae Distribution



### Gibbon



### White handed gibbon



# Gibbon




### Siamang






### Siamang



# The Great Apes

- Families: Pongidae and Hominidae
- Who goes in which?




# Typical Taxonomy

- Pongidae: Orangutan, Gorilla, Chimp, Bonobo
- Hominidae: Humans
- DOES NOT MATCH UNDERSTANDING OF PHYLOGENY

## Modern Taxonomy


- Pongidae = Orangutans
- Hominidae = Gorillas, Chimpanzee, Bonobo, Human
  - Subfamily Gorillinae = gorillas
  - Subfamily Paninae = chimpanzee, bonobo
  - Subfamily Homininae = humans + ancestors



37

# Orangutan Distribution










## African Ape Distribution



### Gorilla




#### Gorilla



#### Gorilla



#### Chimpanzee





46

### Chimpanzee






## Chimp/Bonobo Tool Use





#### Bonobo



#### Bonobo



#### Bonobo



